



Effects of radiofrequency electromagnetic radiation (RF-EMF) on honey bee queen development and mating success

Richard Odemer ^{*}, Franziska Odemer

University of Hohenheim, Apicultural State Institute, 70593 Stuttgart, Germany




HIGHLIGHTS

- Chronic RF-EMF exposure significantly reduced hatching of honey bee queens.
- Mortalities occurred during pupation, not at the larval stages.
- Mating success was not adversely affected by the irradiation.
- After the exposure, surviving queens were able to establish intact colonies.

GRAPHICAL ABSTRACT

Illustrations after Gullan and Cranston (2014)



ARTICLE INFO

Article history:

Received 2 October 2018
 Received in revised form 9 January 2019
 Accepted 13 January 2019
 Available online 15 January 2019

Editor: Henner Hollert

Keywords:

RF-EMF exposure
 Ontogenetic queen development
 Honey bees
Apis mellifera
 Mobile phone radiation
 GSM

ABSTRACT

Mobile phones can be found almost everywhere across the globe, upholding a direct point-to-point connection between the device and the broadcast tower. The emission of radiofrequency electromagnetic fields (RF-EMF) puts the surrounding environment inevitably into contact with this radiation. We have therefore exposed honey bee queen larvae to the radiation of a common mobile phone device (GSM band at 900 MHz) during all stages of their pre-adult development including pupation. After 14 days of exposure, hatching of adult queens was assessed and mating success after further 11 days, respectively. Moreover, full colonies were established of five of the untreated and four of the treated queens to contrast population dynamics. We found that mobile phone radiation had significantly reduced the hatching ratio but not the mating success. If treated queens had successfully mated, colony development was not adversely affected. We provide evidence that mobile phone radiation may alter pupal development, once succeeded this point, no further impairment has manifested in adulthood. Our results are discussed against the background of long-lasting consequences for colony performance and the possible implication on periodic colony losses.

© 2019 Elsevier B.V. All rights reserved.

1. Introduction

The modern world turns around technological achievements and it is simply not possible to imagine our everyday life without them.

^{*} Corresponding author.

E-mail address: richard.odemer@uni-hohenheim.de (R. Odemer).

With an estimated 6.9 billion subscriptions globally, mobile phone devices such as smart phones have established their position in our society (WHO, 2014). In many countries, cell phones are important tools not only for communication but also for bank transfers, newscast, social media and numerous other conveniences with an increasing tendency. Provided that this market will be further growing in the future, concerns are rising about the emission of radiofrequency electromagnetic fields

(RF-EMF) from these devices and their broadcasting network, i.e. antennas and base stations, perceived as environmental pollution (Balmori, 2015).

Radiofrequency waves are electromagnetic fields, and unlike ionizing radiation such as X-rays or gamma rays, they can neither break chemical bonds nor cause ionization in the living tissue (Genuis and Lipp, 2012). They are usually ranging from 30 kHz–300 GHz with cell phones operating mainly between 800 MHz and 3 GHz, pulsed at low frequencies (Hardell, 2017). As a consequence, they are often strictly forbidden in medical facilities and on airplanes, as the radiofrequency signals may interfere with certain electro-medical devices and navigation systems.

In the last decade field and laboratory studies have furthermore demonstrated that RF-EMF exposure is of ecological relevance. The radiation may have an impact on surrounding flora as well as vertebrate and invertebrate organisms (Cucurachi et al., 2013). Effects have manifested in different ways and some of them are a cause of concern. A large scale monitoring study (>10 years) revealed that in trees, a closer range to phone masts resulted in significant damages in the side facing the mast in contrast to the opposite side (Waldmann-Selsam et al., 2016) whereas Roux et al. (2006, 2008) found exposed tomato plants to show similar consequences when wounded, trimmed or burnt. In chicken eggs, Batellier et al. (2008) found an increased mortality when exposed to cell phone radiation over the entire incubation period. Very similar to previous study results from Bastide et al. (2001) and Grigoryev (2003), this developmental stage seems to be particularly vulnerable for non-thermal levels of radiation. A proportional relationship between the intensity of the electromagnetic field and the negative effects, however, could not be established (Batellier et al., 2008).

In fruit flies, reproduction and reproductive organs were also significantly affected by mobile phone radiation (Panagopoulos et al., 2004; Panagopoulos, 2012) unlike to the findings of Weisbrot et al. (2003) where a beneficial effect on the reproductive success was reported. In their study, the number of offspring increased by up to 50% compared to control, demonstrating controversial outcomes. Studies in insects have shown that reproduction cycles and change of generations are quick, making this test system suitable for the detection of possible consequences of RF-EMF exposure. Important biological endpoints such as fertility, reproduction, behavior and development are rather easy to implement, especially in a laboratory setting.

Besides the fruit fly as model organism, special ecological relevance is outlined by pollinators, in particular by the honey bee *Apis mellifera*. They provide critical pollination services valued at over \$200 billion worldwide (Lautenbach et al., 2012), representing 9.5% of the total human food production (Gallai et al., 2009). Even though the honey bee does not seem capable of maximizing agricultural production by replacing ecosystem service of wild pollinators being more efficient - wild insects and honey bees contribute additive to pollination of crops and are, therefore, necessary to optimize the global agricultural pollination service (Garibaldi et al., 2013).

In the past, honey bees have suffered periodic losses within the last century, and in the US a phenomenon called “Colony Collapse Disorder” (CCD) made headlines in the first decade of the new millennium (vanEngelsdorp et al., 2009). Several causative factors have been outlined in the past, among others, pathogens, malnutrition, management, and pesticides have been narrowly focused as main culprits (Steinhauer et al., 2018). One of the most important and economically relevant factors for colony declines worldwide is the predatory mite *Varroa destructor*, introduced in the western world between the late 70ies and 80ies. Since the mites’ global spread, beekeepers are obliged to treat against this parasite, otherwise untreated colonies collapse within a short period of time (Rosenkranz et al., 2010). With reference to CCD, many other factors were also considered to have an impact on honey bee health, however with a rather insignificant regard. A few to name are air pollution (Girling et al., 2013; McFrederick et al., 2008), nanomaterials (Milivojević et al., 2015), solar radiation (Ferrari, 2014),

robbing insects (Core et al., 2012) and global warming (Le Conte and Navajas, 2008).

Worthy of mention, in 2007 a story in an UK newspaper brought to the fore that CCD can be linked to RF-EMF with drastic consequences for bee behavior and homing success (Kimmel et al., 2007; Carreck, 2014). Subsequent studies seem to provide supporting evidence of impaired behavior (Favre, 2011) and affected homing ability (Ferrari, 2014), bearing a potential risk to other bee species such as bumblebees (*Bombus terrestris*) when interacting with floral electric fields and electric field sensing as important sensory modality (Clarke et al., 2013).

However, there are far too few scientific publications to draw a clear conclusion in regard if and to which extent mobile phone radiation represents a real threat to honey bees. A current review actually goes as far as stating that all examined studies were characterized by substantial shortcomings which were sometimes even admitted by their authors upfront, especially in terms of inaccurate dosimetry of mobile phone devices (Verschaeve, 2014). In contrast, Panagopoulos et al. (2016) argued that precisely because the radiation is of highly varying nature, regardless of any dosimetry, radiation levels of mobile phones represent the most realistic exposure.

For a honey bee colony, health and productivity is directly linked to its queen. She represents the growth potential expressed as productivity, being the only egg layer in the collective and therefore responsible for a positive turnover of workers to increase in size at the beginning of each bee season (Moore et al., 2015). In an US survey of winter colony losses, the fourth most important factor identified was due to queen failure (vanEngelsdorp et al., 2008). Given the importance of this individual, our experiments therefore strictly focused on ontogenetic development and further mating success of young queens. We have created a worst case scenario, where mobile phone radiation was adopted by natural means of human exposure. To our knowledge this is the first study that analyzes the effect of a chronic application of mobile phone radiation on honey bee queens. We wanted to prove (i) if under field conditions and good apicultural practice the radiation has any effect at all and to what extent, in addition (ii) we wanted to follow queens which developed under chronic RF-EMF exposure to assess potential risks for the bee colony.

2. Materials & methods

2.1. Field sites and weather conditions

The field sites were located near the Apicultural State Institute in Stuttgart-Hohenheim, Southern Germany (48°42′31.8″N 9°12′38.2″E). At the time present, natural food sources consisted mainly of nectar from diverse local flora such as *Taraxacum officinale*, *Rubus section*, *Tilia* spp. and others. The average temperature during the experiment ranged from 15.2 to 20.1 °C with a precipitation of 90 to 45 L/m². Overall, good weather conditions prevailed for both, mating and foraging (Deutscher Wetterdienst, 2018).

2.2. Experimental setup

This study was performed from May until August in 2018 with healthy queenright colonies from the stock of our apiary. Two replications were employed simultaneously, consisting of two collector colonies: Rep1 (Control1 + EMF1) and Rep2 (Control2 + EMF2). For both approaches, one brood frame with almost fully covered areas of sealed brood and attached bees from eight random colonies were taken out on D-9 and placed in a new ten-frame box, respectively. This box was supplied with two frames of food, as well as a second box on top with ten food frames to ensure sustenance and sufficient room for the hatching bees. Nine days after this procedure (D0), the hive was inspected, and where appropriate, superseded cells were removed to prevent the introduction of a young queen. Further, 18 frames then were split homogeneously but random into two boxes with nine frames each,

complemented with a grafting frame in the center. L1 larvae from a selected colony were grafted and introduced, respectively. Again, grafting of the larvae was randomized by using both sides of the brood comb (A and B). Per replication, 26 larvae (13 A, 13 B) were assigned to each treatment, i.e. control and EMF.

The two boxes then were placed at a different location in approximately 3 km distance to prevent worker bees to return to their original position. Subsequently, at different intervals, assessments were performed to check the no. of accepted larvae after grafting (D1), to protect the capped cells before hatching (D10), to check the hatching rate (D13) and the mating success (D24). After the young queens have hatched, they were transferred to mating units consisting of one of the former brood frames with approximately 1000 bees attached and one food comb.

Successful mating was confirmed on D24 by the presence of eggs, young larvae and capped brood and queens from each treatment (five from the control, four from the treatment) were re-accommodated in new 10-frame boxes to develop into full colonies. After approximately twelve weeks (D88), a colony assessment was performed to record the number of bees and brood. See Fig. 1 for a detailed timeline.

2.3. Mobile phone exposure and measurement

Queen larvae/pupae were treated with a mobile phone (AEG M1220, GSM quad band, China) attached to the grafting frame holding 26 queen cups (Nicot, NICOTPLAST SAS, Maisod, France). The mobile phone handset had a measured specific energy absorption rate (SAR) of 0.59 W/kg (head) and 1.16 W/kg (body worn) according to the international standard IEC 62209-2:2010 (measured with SAR Meter ESM 120, Maschek, Germany - measuring range 0.001–20 W/kg; accuracy 1 mW/kg). Both SAR values were below the 2 W/kg maximum upper limits recommended by the International Commission on Non-Ionizing Radiation Protection (ICNIRP).

A 900 MHz GSM roaming band was used (Vodafone prepaid rate, Vodafone GmbH, Germany). To verify an adequate EMF output, the narrowband radiation was measured three times in three different positions/distances to the mobile phone with a mobile radio frequency

exposure meter to assess electric field strength from 80 MHz up to 3 GHz illustrated in Fig. 2 (ExpoM – RF, frequency band GSM 900 downlink 925–960 MHz, dynamic range 0.005–5 V/m, Fields at Work GmbH, Switzerland). All measurements were carried out in the field, at the same location the experiment was conducted to assure similar exposure conditions. For the measurements we adapted methods proposed by Vermeeren et al. (2013) and Bürgi et al. (2010).

To ensure power supply, the phone was equipped with a power bank (PLOCHY 24,000 mAh Solar, China), the battery status was frequently checked. After the larvae were grafted into the cups by using an appropriate tool, 15 telephone calls with a two minute duration were applied every 24 h for a total of two weeks (non-speaking emission, GSM 900 downlink) at random equivalent to a daily operator exposure of 30 min. The device was turned off in the control group for sham exposure.

2.4. Colony assessment

The amount of bees and brood cells (open and sealed) were estimated with the Liebefeld Method (Imdorf et al., 1987), which is a feasible tool to provide accurate and reliable evaluation of colony strength (measuring error $\pm 10\%$). Care was taken that all colonies were evaluated by the same person to minimize variation and colony assessment was conducted in the morning before bee flight.

2.5. Statistical analysis

We evaluated the mortality data with a Kaplan-Meier-Survival analysis (KM). Survivorship between control and treatment was compared pairwise and tested for significance with a Log-Rank Tests (Cox-Mantel). Individuals collected at the end of the experiment were considered censored, as were those observed but not collected on the final day. Furthermore, larvae that disappeared during the experiment were considered dead on the last day they were seen. Both treatment groups and the two replicates (Rep1 = Control1 + EMF1; Rep2 = Control2 + EMF2) were additionally compared with a Cox proportional hazards model to determine the hazard ratio (HR). Possible inter-colony effects




Fig. 1. Timeline of the experiment. At D-9, eight brood frames with attached bees were taken out from respective donor colonies and placed in one collector colony. At D0, supersedure cells were removed and the collector colony was split in two sub-colonies. In addition, a grafting frame with L1 larvae was inserted. RF-EMF exposure lasted until D13, when queens were about to hatch. Young queens were subsequently inserted into mating units where mating success was checked at D24. Successfully mated queens with one frame of approximately 1000 bees were relocated into new boxes where they were able to establish a new colony. Finally, at D88 the condition of these colonies was assessed.




Fig. 2. Grafting frame placed in the EMF treatment colony containing 26 queen cups. The mobile phone device was attached in the center of the frame, its radiation intensity is indicated with the differently colored sections in the illustration (darker > lighter). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

were evaluated as covariate to justify pooling data of the same treatments. In addition, both control groups and both EMF groups were statistically compared to further confirm pooling of these datasets with KM and HR. The estimated number of bees and brood cells were checked with a Shapiro-Wilk test for normal distribution. If data was normal, a one-way ANOVA was performed on the two experimental groups, respectively. For all tests RStudio (R Core Team, 2018) and significance level of $\alpha = 0.05$ was used.

3. Results

3.1. Honey bee queen survival

The Kaplan-Meier-Survival analysis of both groups showed a significant difference indicating a higher mortality of the EMF treated bees when compared to the control group ($p = 0.0054$) (Fig. 3). In addition, a Cox proportional hazards model was applied to determine the hazard ratio (HR) displayed as forest plot (Fig. 4). With a HR of 2.3 the EMF treated queens had a significantly increased risk of dying when

compared to the control ($p = 0.003$). Moreover, the two replicates (Rep1 and Rep2) were compared as covariate to display possible inter-colony effects. However, with a HR of 1.7 queens in Rep2 did not have a higher risk of dying when compared to Rep1 ($p = 0.062$), therefore data of both replicates were pooled. More details are displayed in the supplementary figure Figs. S1 to S4.

3.2. Hatching and mating success

The acceptance rate of grafted larvae on D1 was 76.9% and identical in both treatments. As shown in Fig. 3, a significant decrease of individuals in the EMF treatment occurred during the pupation phase of the experiment. At D10, queen cells were protected with a plastic cage to prevent hatching queens from killing each other. We observed a similar decrease of 26.9 (control) to 34.6% (treatment) at this stage compared to the initially grafted cells on D0 (Log-Rank $p = 0.43$), being not significantly different. The hatch of adult queens at D13 revealed a significant decrease of formerly treated queens during pupation when compared to D0 with a reduction of 30.8% in the control to 61.5% in the treatment




Fig. 3. Both groups were compared with a Kaplan-Meier-Survival analysis. A post hoc Log-Rank test (Cox-Mantel) revealed a significant higher mortality in the EMF treatment when compared to the control (Log-Rank $p = 0.0054$), where a significant decrease of individuals occurred during the pupation phase of the experiment (see also Fig. 5).




Fig. 4. Both treatment groups and the two replicates (Rep1 and Rep2) were additionally compared with a Cox proportional hazards model to determine the hazard ratio (HR) displayed as forest plot. With a HR of 2.3 the EMF treated queens had a significantly increased risk of dying when compared to the control ($p = 0.003$). And with a HR of 1.7 queens in Rep2 did not have a higher risk of dying when compared to Rep1 ($p = 0.062$).

(Log-Rank $p = 0.0065$), representing a difference of 44.4% between both groups. Mating success (D24) was compared to the total of hatched queens on D13 showing no significant changes (Log-Rank $p = 0.18$) (see also the supplementary figures Figs. S5 to S10).

3.3. Colony assessment

The population of bees and brood cells was estimated at D88. The results are shown in Fig. 6A for the number of bees and in Fig. 6B for the number of brood cells. We compared the two treatment groups with a

one-way ANOVA but could not see significant differences for the number of bees ($p = 0.688$) or the amount of brood cells ($p = 0.768$).

4. Discussion

The emission of radiofrequency electromagnetic fields (RF-EMF) and their negative effects towards honey bee health has been controversially discussed in the past (Carreck, 2014; Verschaeve, 2014; Panagopoulos et al., 2016). Here, we could demonstrate for the first time that RF-EMF exposure has significantly affected ontogenetic




Fig. 5. Total number of L1 larvae grafted and followed through their ontogenetic development from pupa to adult. In the EMF treatment a significant decrease of individuals came into effect within the pupation phase of the experiment (*; Log-Rank $p = 0.0065$) but the mating success was not affected (ns; Log-Rank $p = 0.18$, see also Figs. 3 and S5 to S10). Illustrations after Gullan and Cranston (2014).




Fig. 6. A. Number of bees estimated at D88 in the colonies of the control ($n = 5$) and of the EMF treatment ($n = 4$). Same letters indicate no statistically significant differences ($p = 0.688$, ANOVA). B. Number of brood cells estimated at D88 in the colonies of the control ($n = 5$) and of the EMF treatment ($n = 4$). Same letters indicate no statistically significant differences ($p = 0.768$, ANOVA).

queen development under worst case operator conditions. We observed an increased mortality during pupation resulting in a reduced hatching rate of the later queens. This is in line with a reduced reproductive capacity found in fruit flies (*Drosophila melanogaster*) (Panagopoulos et al., 2004; Margaritis et al., 2014), where a linear decrease of fecundity was reported with the duration of exposure (Panagopoulos and Margaritis, 2010). This decrease was further associated with the distance to the mobile phone device showing the strongest effects at <10 cm (Panagopoulos et al., 2010). In our setup, the most distant queen cups were approximately 21 cm away from the radiation source and we therefore assume that for all larvae a worst case scenario came into effect. In addition, the impairment of fruit flies seemingly depended on field intensity (Panagopoulos et al., 2007) not only reducing the offspring but also the ovarian size of the exposed subjects (Panagopoulos, 2012).

At present, only a few studies have investigated the influence of irradiation on insect development. As an example, larvae and

pupae of the dried fruit beetle (*Carpophilus hemipterus*) and the yellow fever mosquito (*Aedes aegypti*) were exposed to Gamma radiation (ionizing radiation). The radiotherapy did not cause acute death in larvae but decreased pupation significantly, no effects however could be observed when either young or old pupae were exposed (Johnson, 1987; Akter and Khan, 2014). It seems likely that RF-EMF had a similar effect in our study, as larval mortality was not elevated. However, this should be further underpinned by exposing larvae and pupae separately. Negative effects should also be specified in more detail by the use of proper biochemical methods to determine oxidative stress or DNA damage as described by Vilić et al. (2017). They found honey bee worker larvae significantly affected when exposed to simulated RF-EMF radiation (voice modulation, 217 Hz), resulting in DNA damage and further corroborating our hypothesis that here rather pupal than larval development was impaired as our device has only emitted radiation in the less bioactive non-speaking mode (Panagopoulos et al., 2004).

Although Hsu et al. (2007) observed changes in the size of magnetic granules in the trophocytes of adult honey bees upon applying a magnetic field to these cells, we could show that mating success remained unaffected suggesting that navigation and the possible disruption of magneto reception or other negative impacts came not into effect, were at least not long-lasting or even reversible (Vacha et al., 2009). Interestingly, we provide evidence that developing honey bee queens once they have survived RF-EMF exposure seem to retain the ability to establish an intact colony. This is indicated by similarly strong numbers of bees and the amount of brood in both our treatment groups with the absence of any signs of impairment (e.g. patchy brood pattern). As a further critical step of colony survival however, overwintering should also be assessed to elucidate possible long term effects from the irradiation (Smart et al., 2016).

The social entity as a whole is able to buffer environmental stressors of various kinds as an expression of social resilience (Straub et al., 2015). Worker bees are nursing eggs and feeding larvae of different casts in their social state, potentially contributing to this mechanism. Here we focused on the development of individual queens from larvae to adult, however, the outcome of our study could also be influenced by the condition of the collector colonies that we have created but not further assessed. Eggs, larvae and pupae are very sensitive stages of development and intensive care is taken to supply their substantial needs in terms of nutrition and environmental conditions, i.e. maintaining a constant temperature and humidity (Wang et al., 2015; Eouzan et al., 2018). RF-EMF radiation is known to affect bees behavior in different ways (Favre, 2011; Ferrari, 2014), which makes it plausible that brood care could also be adversely affected. This important factor should be further investigated and included in future experiments.

With an increasing number of mobile phone devices and as a consequence of good accessibility a higher density of phone masts, not only urban but also rural areas in particular are more and more exposed to irradiation (Balmori, 2009). A measurement of RF-EMF intensities across different European cities revealed maximum field strength values ranging from 0.59 to 0.84 V/m (Urbinello et al., 2014a), respectively, with a maximum value of 0.69 V/m in public transport (Sagar et al., 2016). In contrast, the field strength measured in our study seemed to be beyond these values but in line with Mamrot et al. (2015), demonstrating that the intermittent stress on the test subject (s) can be many fold higher than average levels measured in the surroundings, emitted from generators or found in agglomerations. Our findings confirm that there is a high variability in mobile phone emission (Frei et al., 2009), representing an important feature in terms of bioactivity towards living organism's defense against environmental stressors (Panagopoulos et al., 2015). The authors therefore suggest not using simulated but real mobile phone emissions in an experimental setup, which we have considered. In addition, we have tried to apply a realistic operator exposure scenario in terms of average number of calls and average call duration performed with mobile phone devices. The mobile call duration reported by the German Federal Network Agency (2011) was 2.5 min per call, in Shum et al. (2011) ranging from 2.1 min (self-reports) to 2.8 min (billing records) and <2 min in Friebel and Seabright (2011). Further, the average number of calls per day ranging from 4.1 (Shum et al., 2011) to 5 per day in adults (Lenhart, 2010). In contrast, an average of 33.1 min was reported for total mobile phone call duration from undergraduate college students per day in the US (Roberts et al., 2014). We therefore decided to employ 2 min per call and 15 calls per day resulting in 30 min exposure per day in our experiment, representing a realistic worst case operator exposure.

To achieve a broader understanding how RF-EMF potentially influences the honey bee superorganism, Verschaeve (2014) pointed out that it is mandatory to emphasize particularly the level and duration of exposure, in presence of the relevant environmental situation. Mobile phones, however, use complex algorithms that make it nearly impossible to determine which parameters (intensity,

frequency, modulation, phase, etc.) when and to which degree radiate on a particular test object. This was also demonstrated by the field strength intensities emitted from our mobile phone attached to the grafting frame. Queen cells were not exposed homogeneously, as with distance the radiation got weaker. Regardless of any dosimetry, Panagopoulos et al. (2016) argue for the use of mobile phones as radiation source because it creates a scenario to which billions of operators are exposed every day. We have tried to reproduce a realistic worst case exposure scenario in our experiment, sharing this view. To provide a more precise image of the emitted radiation of mobile phones in future experiments however, software manipulation could be a tool to track the transmitted power densities by the device itself similar to a data logger (Bhatt et al., 2018).

So far, different exposure scenarios were applied in honey bees and a broad range of effects are reported (Cucurachi et al., 2013). Some studies even claimed with RF-EMF to have found the major cause for CCD (Carreck, 2014). However, many of these studies had substantial deficits such as a very low sample size (Sharma and Kumar, 2010), intransparent methods (Sahib, 2011; Kumar et al., 2011; Dalio, 2015) or were even preliminary and did not undergo peer-review (Kimmel et al., 2007). Therefore, findings of this quality were generally not considered reliable in their contribution to colony losses and are far from conclusive (Carreck, 2014).

Since the first reports of "Colony Collapse Disorder" in the US within the first decade of the new millennium, this phenomenon received huge public attention all across the globe. Wherever outside of the US colony losses were reported, a connection to CCD was drawn quickly (Neumann and Carreck, 2010). Considering the described symptoms however, neither colonies in Europe nor on other continents than North America seemed to have expressed them - indicating a high complexity on the local scale of colony declines - making it difficult to narrow down the cause to one single factor. Moreover, periodic high losses of colonies have already occurred throughout the last century, even before mobile phones were invented and used widely (vanEngelsdorp and Meixner, 2009). With the introduction of *V. destructor* a chemical treatment of the colonies became mandatory ensuring their survival (Rosenkranz et al., 2010). Even though the mite has been identified as key for worldwide colony declines, other factors may exert negative influence on bee health, especially if varroosis may have increased their susceptibility. Pesticides, in particular the class of neurotoxic neonicotinoids, received attention after a number of bee poisoning incidents followed by the application of agricultural practices (Rosenkranz et al., 2016) finally leading to a total ban of three neonicotinoid substances in the EU (EFSA, 2018).

As a consequence, political, social and economic awareness to colony declines increased during the last ten years to a significant level in most western civilizations being present up to date in a huge variety of media formats on both, a local and global scale (Potts et al., 2010). Receiving increased public pressure for more information on bee declines, many research projects such as the international COLOSS network (Prevention of honey bee COLony LOSSes, <http://www.coloss.org>) the EPILOBEE consortium (A pan-European epidemiological study on honeybee colony losses, https://ec.europa.eu/food/animals/live_animals/bees/study_on_mortality_en) or the German DEBIMO (Deutsches Bienenmonitoring, <https://bienenmonitoring.uni-hohenheim.de>) were funded and came to fruition. Results of these broad scale monitoring projects provide strong evidence that the survival of a colony depends to a large extent on the skill of the beekeeper as well as on proper disease control (Jacques et al., 2017). Obviously, colonies must reach a population size threshold to survive adverse conditions (Genersch et al., 2010; Döke et al., 2018) which is directly linked to management practices and beekeeper education. Steinhauer et al. (2018) summarized the current leading drivers for colony losses where, besides the above mentioned factors, malnutrition and the use of pesticides are also of high importance. This demonstrates that colony losses are a result of multifactorial interactions between environmental influences and beekeeping

practices (Molineri et al., 2018) continuously challenging bee health to ever-changing factors.

Even though scientific evidence for a contribution of mobile phone radiation to colony declines are currently lacking, we could demonstrate for the first time a clear negative impact on honey bee queens. Taking into consideration that varroosis may affect susceptibility to negative environmental influences, increasing EMF intensities could be weakening the immune system of bees and in return making them more vulnerable to other diseases (Walleczek, 1992; Pall, 2013). This may be of particular importance in view of the current development in the beekeeping sector. As a trend of the last decades, beekeeping became famous with the life style of townsmen all across the globe (Lorenz and Stark, 2015; Kohsaka et al., 2017; Stange et al., 2017). Therefore, density of bee colonies held in urban areas has dramatically increased and may favor the spread of diseases or pathogens (Youngsteadt et al., 2015). However, following this trend also bears the risk of a higher exposure to mobile phone emissions, which seem to be continuously increasing in major cities (Urbanello et al., 2014b), potentially affecting bee health in a future scenario. It might also be worthy to look into parasite-host-interactions of the honey bee, *V. destructor* in particular, where a disturbance through RF-EMF in host-finding could actually be a benefit (Frey et al., 2013).

Gajger et al. (2017) demonstrated weight differences and a lower sperm count in queens treated with a neonicotinoid pesticide. For future experiments with queens exposed to mobile phone radiation these parameters could also be considered to demonstrate a possible negative impact on queen's physiology. In addition, some older studies imply that extremely low-frequency (ELF) electric and magnetic fields emitted by power lines can have local adverse effects on honey bees - mainly due to changes in behavior (stinging). These effects, however, quickly diminish with distance to the power lines or in absence of a conducting medium (Greenberg et al., 1981; Bindokas et al., 1988a, 1988b, 1989). At this current time, not many studies are available that are investigating the influence of such irradiation on bees and other important pollinators. It has even been suggested to create pollinator reservoirs beneath power corridors for an optimal land use and as a benefit for many insects (Russell et al., 2018) without further enquiries or thought about the associated hazards. Yet, it still remains unclear to what extent electromagnetic fields can possibly influence these microenvironments.

5. Conclusion

Even though detrimental effects on ontogenetic queen development were revealed by the outcome of our study, caution is needed in interpreting these results. So far, there have been no serious records of colony losses associated with mobile phone radiation. Moreover, we have created by far a worst case operator scenario to which honey bee colonies would not be exposed under realistic beekeeping conditions. Duration and level were similar to average operator exposure by the use of a mobile phone, but not to those present at an apiary, neither in rural nor in urban areas. And yet, queens that survived the treatment were able to establish full functional colonies, demonstrating an immense recovering potential. Therefore we do not assume any acute negative effects on bee health in the mid-term. However, we do not rule out an influence through lower doses of permanent irradiation, in particular on a chronic sublethal level present in major city environments. Hence, we urgently suggest further research should be carried out in the long-term to ascertain what impacts are to be expected in the context of a suitable risk assessment for electromagnetic fields on bee health.

CRedit authorship contribution statement

Richard Odemer: Conceptualization. **Franziska Odemer:** Conceptualization.

Acknowledgements

Many thanks to Simon Hummel for providing adequate breeding lines to conduct research with extremely calm honey bees and to the anonymous reviewers for substantially improving our manuscript.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declarations of interest

None.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Appendix A. Supplementary figures

Supplementary figures (Figs. S1–S10) to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2019.01.154>.

References

- Akter, H., Khan, S.A., 2014. Sensitivity of immature stages of dengue causing mosquito, *Aedes aegypti* (L.) to gamma radiation. *J. Entomol.* 11, 56–67. <https://doi.org/10.3923/je.2014.56.67>.
- Balmori, A., 2009. Electromagnetic pollution from phone masts. Effects on wildlife. *Pathophysiology* 16, 191–199. <https://doi.org/10.1016/j.pathophys.2009.01.007>.
- Balmori, A., 2015. Anthropogenic radiofrequency electromagnetic fields as an emerging threat to wildlife orientation. *Sci. Total Environ.* 518–519, 58–60. <https://doi.org/10.1016/j.scitotenv.2015.02.077>.
- Bastide, M., Youbicier-Simo, B.J., Lebecq, J.C., Giannis, J., 2001. Toxicological study of electromagnetic radiation emitted by television and video display screens and cellular telephones on chickens and mice. *Indoor Built Environ.* 10, 291–298.
- Batellier, F., Couty, I., Picard, D., Brillard, J.P., 2008. Effects of exposing chicken eggs to a cell phone in 'call' position over the entire incubation period. *Theriogenology* 69, 737–745.
- Bhatt, C.R., Redmayne, M., Abramson, M.J., Sim, M.R., Brzozek, C., Zeleke, B.M., Benke, G., 2018. Estimating transmitted power density from mobile phone: an epidemiological pilot study with a software modified phone. *Australas. Phys. Eng. Sci. Med.* 41, 985–991. <https://doi.org/10.1007/s13246-018-0699-7>.
- Bindokas, V.P., Gauger, J.R., Greenberg, B., 1988a. Exposure scheme separates effects of electric shock and electric field for honey bees, *Apis mellifera* L. *Bioelectromagnetics* 9, 275–284.
- Bindokas, V.P., Gauger, J.R., Greenberg, B., 1988b. Mechanism of biological effects observed in honey bees (*Apis mellifera*, L.) hived under extra-high-voltage transmission lines: implications derived from bee exposure to simulated intense electric fields and shocks. *Bioelectromagnetics* 9, 285–301.
- Bindokas, V.P., Gauger, J.R., Greenberg, B., 1989. Laboratory investigations of the electrical characteristics of honey bees and their exposure to intense electric fields. *Bioelectromagnetics* 10, 1–12.
- Bürgi, A., Frei, P., Theis, G., Mohler, E., Braun-Fahrlander, C., Fröhlich, J., Neubauer, G., Egger, M., Rössli, M., 2010. A model for radiofrequency electromagnetic field predictions at outdoor and indoor locations in the context of epidemiological research. *Bioelectromagnetics* 31, 226–236. <https://doi.org/10.1002/bem.20552>.
- Carreck, N., 2014. Electromagnetic radiation and bees, again. *Bee World* 91 (4), 101–102. <https://doi.org/10.1080/0005772X.2014.11417624>.
- Clarke, D., Whitney, H., Sutton, G., Robert, D., 2013. Detection and learning of floral electric fields by bumblebees. *Science* (80-) 340, 66–69. <https://doi.org/10.1126/science.1230883>.
- Core, A., Runckel, C., Ivers, J., Quock, C., Siapno, T., DeNault, S., Brown, B., DeRisi, J., Smith, C.D., Hafernik, J., 2012. A new threat to honey bees, the parasitic phorid fly *Apocephalus borealis*. *PLoS One* 7, e29639. <https://doi.org/10.1371/journal.pone.0029639>.
- Cucurachi, S., Tamis, W.L.M., Vijver, M.G., Peijnenburg, W.J.G.M., Bolte, J.F.B., de Snoo, G.R., 2013. A review of the ecological effects of radiofrequency electromagnetic fields (RF-EMF). *Environ. Int.* 51, 116–140. <https://doi.org/10.1016/j.envint.2012.10.009>.
- Dalio, J.S., 2015. Effect of electromagnetic (cell phone) radiations on *Apis mellifera*. *J. Res. Agric. Anim. Sci.* 2 (12), 06–10.
- Deutscher Wetterdienst, D.W.D., 2018. Deutschlandwetter im Sommer 2018. Pressemitteilung. <http://www.dwd.de>, Accessed date: 26 September 2018.
- Döke, M.A., McGrady, C.M., Otieno, M., Grozinger, C.M., Frazier, M., 2018. Colony size, rather than geographic origin of stocks, predicts overwintering success in honey bees (Hymenoptera: Apidae) in the northeastern United States. *J. Econ. Entomol.* 1–9. <https://doi.org/10.1093/jee/toy377>.
- EFSA, 2018. Evaluation of the data on clothianidin, imidacloprid and thiamethoxam for the updated risk assessment to bees for seed treatments and granules in the EU. EFSA Support. Publ. 15, 1–31. <https://doi.org/10.2903/sp.efsa.2018.EN-1378>.

- Eouzan, I., Garnery, L., Pinto, M.A., Delalande, D., Neves, C.J., 2018. Hygroregulation, a key ability for eusocial insects: Native Western European bees as a case study. *bioRxiv* <https://doi.org/10.1101/351874>.
- Favre, D., 2011. Mobile phone-induced honeybee worker piping. *Apidologie* 42, 270–279. <https://doi.org/10.1007/s13592-011-0016-x>.
- Federal Network Agency, 2011. Press release 28 July 2011: average duration of 2.5 minutes for mobile calls. Bonn, Germany. <http://www.bundesnetzagentur.de>, Accessed date: 24 August 2018.
- Ferrari, T.E., 2014. Magnets, magnetic field fluctuations and geomagnetic disturbances impair the homing ability of honey bees (*Apis mellifera*). *J. Apic. Res.* 53, 452–465. <https://doi.org/10.3896/IBRA.1.53.4.15>.
- Frey, P., Mohler, E., Neubauer, G., Theis, G., Bürgi, A., Fröhlich, J., Braun-Fahrlander, C., Bolte, J., Egger, M., Rössli, M., 2009. Temporal and spatial variability of personal exposure to radio frequency electromagnetic fields. *Environ. Res.* 109, 779–785. <https://doi.org/10.1016/j.envres.2009.04.015>.
- Frey, E., Odemer, R., Blum, T., Rosenkranz, P., 2013. Activation and interruption of the reproduction of *Varroa destructor* is triggered by host signals (*Apis mellifera*). *J. Invertebr. Pathol.* 113, 56–62. <https://doi.org/10.1016/j.jip.2013.01.007>.
- Friebel, G., Seabright, P., 2011. Do women have longer conversations? Telephone evidence of gendered communication strategies. *J. Econ. Psychol.* 32, 348–356. <https://doi.org/10.1016/j.joep.2010.12.008>.
- Gajger, I.T., Sakač, M., Gregorc, A., 2017. Impact of thiamethoxam on honey bee queen (*Apis mellifera carnica*) reproductive morphology and physiology. *Bull. Environ. Contam. Toxicol.* 99, 297–302. <https://doi.org/10.1007/s00128-017-2144-0>.
- Gallai, N., Salles, J.M., Settele, J., Vaissière, B.E., 2009. Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecol. Econ.* 68, 810–821. <https://doi.org/10.1016/j.ecolecon.2008.06.014>.
- Garibaldi, L.A., Steffan-Dewenter, I., Winfree, R., Aizen, M.A., Bommarco, R., Cunningham, S.A., Kremen, C., Carvalheiro, L.G., Harder, L.D., Afik, O., Bartomeus, I., Benjamin, F., Boreux, V., Cariveau, D., Chacoff, N.P., Dudenhofer, J.H., Freitas, B.M., Ghazoul, J., Greenleaf, S., Hipolito, J., Holzschuh, A., Howlett, B., Isaacs, R., Javorek, S.K., Kennedy, C.M., Krewenka, K.M., Krishnan, S., Mandelik, Y., Mayfield, M.M., Motzke, I., Munyuli, T., Nault, B.A., Otieno, M., Petersen, J., Pisanty, G., Potts, S.G., Rader, R., Ricketts, T.H., Rundlof, M., Seymour, C.L., Schuepp, C., Szentgyorgyi, H., Taki, H., Tscharrnke, T., Vergara, C.H., Viana, B.F., Wanger, T.C., Westphal, C., Williams, N., Klein, A.M., 2013. Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science* (80-) 339, 1608–1611. <https://doi.org/10.1126/science.1230200>.
- Genersch, E., Von Der Ohe, W., Kaatz, H., Schroeder, A., Otten, C., Büchler, R., Berg, S., Ritter, W., Mühlen, W., Gisdler, S., Meixner, M., Liebig, G., Rosenkranz, P., 2010. The German bee monitoring project: a long term study to understand periodically high winter losses of honey bee colonies. *Apidologie* 41, 332–352. <https://doi.org/10.1051/apido/2010014>.
- Genius, S.J., Lipp, C.T., 2012. Electromagnetic hypersensitivity: fact or fiction? *Sci. Total Environ.* 414, 103–112. <https://doi.org/10.1016/j.scitotenv.2011.11.008>.
- Girling, R.D., Lusebrink, I., Farthing, E., Newman, T.A., Poppy, G.M., 2013. Diesel exhaust rapidly degrades floral odours used by honeybees. *Sci. Rep.* 3, 2779. <https://doi.org/10.1038/srep02779>.
- Greenberg, B., Bindokas, V.P., Gauger, J.R., 1981. Biological effects of 765-kV transmission line: exposures and thresholds in honeybee colonies. *Bioelectromagnetics* 2, 315–328.
- Grigoryev, Y., 2003. Biological effects of mobile phone electromagnetic field on chick embryo (risk assessment using the mortality rate). *Radiats. Biol. Radioecol.* 43, 541–543.
- Gullan, P.J., Cranston, P.S., 2014. *The Insects: An Outline of Entomology*. Fifth edition. John Wiley & Sons Blackwell, Oxford 1-4051-1113-5.
- Hardell, L., 2017. World Health Organization, radiofrequency radiation and health - a hard nut to crack (review). *Int. J. Oncol.* 51, 405–413. <https://doi.org/10.3892/ijo.2017.4046>.
- Hsu, C.-Y., Ko, F.-Y., Li, C.-W., Fann, K., Lue, J.-T., 2007. Magnetoreception system in honeybees (*Apis mellifera*). *PLoS One* 2, e395. <https://doi.org/10.1371/journal.pone.0000395>.
- Imdorf, A., Buehlmann, G., Gerig, L., Kilchenmann, V., 1987. Überprüfung der Schätzmethode zur Ermittlung der Brutfläche und der Anzahl Arbeiterinnen in freifliegenden Bienenvölkern. *Apidologie* 18, 137–146. <https://doi.org/10.1051/apido:19870204>.
- Jacques, A., Laurent, M., Ribière-Chabert, M., Saussac, M., Bougeard, S., Budge, G.E., Hendriks, P., Chauzat, M.-P., 2017. A pan-European epidemiological study reveals honey bee colony survival depends on beekeeper education and disease control. *PLoS One* 12, e0172591. <https://doi.org/10.1371/journal.pone.0172591>.
- Johnson, J.A., 1987. Sensitivity of larvae, pupae, and adults of the driedfruit beetle (*Coleoptera: Nitidulidae*) to gamma radiation. *J. Econ. Entomol.* 80, 1302–1305.
- Kimmel, S., Kuhn, J., Harst, W., Stever, H., 2007. Electromagnetic Radiation: Influences on Honeybees (*Apis mellifera*). unpublished. pp. 1–6.
- Kohsaka, R., Park, M.S., Uchiyama, Y., 2017. Beekeeping and honey production in Japan and South Korea: past and present. *J. Ethn. Foods* 4, 72–79. <https://doi.org/10.1016/j.jef.2017.05.002>.
- Kumar, N.R., Sangwan, S., Badotra, P., 2011. Exposure to cell phone radiations produces biochemical changes in worker honey bees. *Toxicol. Int.* 18 (1), 70–72.
- Lautenbach, S., Seppelt, R., Liebscher, J., Dormann, C.F., 2012. Spatial and temporal trends of global pollination benefit. *PLoS One* <https://doi.org/10.1371/journal.pone.0035954>.
- Le Conte, Y., Navajas, M., 2008. Climate change: impact on honey bee populations and diseases. *Rev. Sci. Tech.* 27 (485–97), 499–510.
- Lenhart, A., 2010. Cell phones and American adults: they make just as many calls, but text less often than teens. *Pew Res. Cent.* 1–42.
- Lorenz, S., Stark, K., 2015. Saving the honeybees in Berlin? A case study of the urban bee-keeping boom. *Environ. Sociol.* 1 (2), 116–126. <https://doi.org/10.1080/23251042.2015.1008383>.
- Mamrot, P., Mariańska, M., Aniolczyk, H., Politański, P., 2015. Electromagnetic fields in the vicinity of DECT cordless telephones and mobile phones. *Med. Pr.* 66, 803–814. <https://doi.org/10.13075/mp.5893.00364>.
- Margaritis, L.H., Manta, A.K., Kokkialaris, K.D., Schiza, D., Alimisis, K., Barkas, G., Georgiou, E., Giannakopoulou, O., Kollia, I., Kontogianni, G., Kourouzidou, A., Myari, A., Roumelioti, F., Skouroliakou, A., Sykioti, V., Varda, G., Xenos, K., Ziomias, K., 2014. *Drosophila* oogenesis as a bio-marker responding to EMF sources. *Electromagn. Biol. Med.* 33, 165–189. <https://doi.org/10.3109/15368378.2013.800102>.
- McFrederick, Q.S., Kathilankal, J.C., Fuentes, J.D., 2008. Air pollution modifies floral scent trails. *Atmos. Environ.* 42, 2336–2348. <https://doi.org/10.1016/j.atmosenv.2007.12.033>.
- Milivojević, T., Glavan, G., Božič, J., Sepčić, K., Mesarič, T., Drobne, D., 2015. Neurotoxic potential of ingested ZnO nanomaterials on bees. *Chemosphere* 120, 547–554. <https://doi.org/10.1016/j.chemosphere.2014.07.054>.
- Molineri, A., Giacobino, A., Pacini, A., Bulacio Cagnolo, N., Merke, J., Orellano, E., Bertozzi, E., Zago, L., Aignasse, A., Pietronave, H., Rodríguez, G., Crisanti, P., Palacio, M.A., Signorini, M., 2018. Environment and *Varroa destructor* management as determinant of colony losses in apiaries under temperate and subtropical climate. *J. Apic. Res.* 57, 551–564. <https://doi.org/10.1080/00218839.2018.1475697>.
- Moore, P.A., Wilson, M.E., Skinner, J.A., 2015. Department of Entomology and Plant Pathology. The University of Tennessee, Knoxville TN <https://bit.ly/2APXpSx>, Accessed date: 24 September 2018.
- Neumann, P., Carreck, N.L., 2010. Honey bee colony losses. *J. Apic. Res.* 49 (1–6). <https://doi.org/10.3896/IBRA.1.49.1.01>.
- Pall, M.L., 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J. Cell. Mol. Med.* 17, 958–965. <https://doi.org/10.1111/jcmm.12088>.
- Panagopoulos, D.J., 2012. Effect of microwave exposure on the ovarian development of *Drosophila melanogaster*. *Cell Biochem. Biophys.* 63, 121–132. <https://doi.org/10.1007/s12013-012-9347-0>.
- Panagopoulos, D.J., Margaritis, L.H., 2010. The effect of exposure duration on the biological activity of mobile telephony radiation. *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* 699, 17–22. <https://doi.org/10.1016/j.mrgentox.2010.04.010>.
- Panagopoulos, D.J., Karabarbounis, A., Margaritis, L.H., 2004. Effect of GSM 900-MHz mobile phone radiation on the reproductive capacity of *Drosophila melanogaster*. *Electromagn. Biol. Med.* 23, 29–43. <https://doi.org/10.1081/JBC-120039350>.
- Panagopoulos, D.J., Chavdoula, E.D., Karabarbounis, A., Margaritis, L.H., 2007. Comparison of bioactivity between GSM 900 MHz and DCS 1800 MHz mobile telephony radiation. *Electromagn. Biol. Med.* 26, 33–44. <https://doi.org/10.1080/15368370701205644>.
- Panagopoulos, D.J., Chavdoula, E.D., Margaritis, L.H., 2010. Bioeffects of mobile telephony radiation in relation to its intensity or distance from the antenna. *Int. J. Radiat. Biol.* 86, 345–357. <https://doi.org/10.3109/09553000903567961>.
- Panagopoulos, D.J., Johansson, O., Carlo, G.L., 2015. Real versus simulated mobile phone exposures in experimental studies. *Biomed. Res. Int.* <https://doi.org/10.1155/2015/607053>.
- Panagopoulos, D.J., Cammaerts, M.-C., Favre, D., Balmori, A., 2016. Comments on environmental impact of radiofrequency fields from mobile phone base stations. *Crit. Rev. Environ. Sci. Technol.* 46, 885–903. <https://doi.org/10.1080/10643389.2016.1182107>.
- Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O., Kunin, W.E., 2010. Global pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* 25, 345–353. <https://doi.org/10.1016/j.tree.2010.01.007>.
- R Core Team, 2018. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Roberts, J., Yaya, L., Manolis, C., 2014. The invisible addiction: cell-phone activities and addiction among male and female college students. *J. Behav. Addict.* 3, 254–265. <https://doi.org/10.1556/JBA.3.2014.015>.
- Rosenkranz, P., Aumeier, P., Ziegelmann, B., 2010. Biology and control of *Varroa destructor*. *J. Invertebr. Pathol.* 103 (Suppl.), S96–119. <https://doi.org/10.1016/j.jip.2009.07.016>.
- Rosenkranz, P., von der Ohe, W., Schäfer, M., Genersch, E., Büchler, R., Berg, S., Otten, C., 2016. Veröffentlichungen und Berichte Website Deutsches Bienenmonitoring – “DeBiMo”. <https://goo.gl/JBj8LC>, Accessed date: 8 January 2019.
- Roux, D., Vian, A., Girard, S., Bonnet, P., Paladian, F., Davies, E., Ledoigt, G., 2006. Electromagnetic fields (900 MHz) evoke consistent molecular responses in tomato plants. *Physiol. Plant.* 128, 283–288.
- Roux, D., Vian, A., Girard, S., Bonnet, P., Paladian, F., Davies, E., Ledoigt, G., 2008. High frequency (900 MHz) low amplitude (5 V m^{-1}) electromagnetic field: a genuine environmental stimulus that affects transcription, translation, calcium and energy charge in tomato. *Planta* 227, 883–891.
- Russell, K.N., Russell, G.J., Kaplan, K.L., Mian, S., Kornbluth, S., 2018. Increasing the conservation value of powerline corridors for wild bees through vegetation management: an experimental approach. *Biodivers. Conserv.* 27, 2541–2565. <https://doi.org/10.1007/s10531-018-1552-8>.
- Sagar, S., Struchen, B., Finta, V., Eeftens, M., Rössli, M., 2016. Use of portable exposimeters to monitor radiofrequency electromagnetic field exposure in the everyday environment. *Environ. Res.* 150, 289–298. <https://doi.org/10.1016/j.envres.2016.06.020>.
- Sahib, S., 2011. Electromagnetic radiation (EMR) clashes with honey bees. *Int. J. Environ. Sci.* 1, 897–900.
- Sharma, M.C., Kumar, N., 2010. Changes in honeybee behaviour and biology under the influence of cellphone radiations. *Vet. Parasitol.* 154, 98–102. <https://doi.org/10.1016/j.vetpar.2008.03.001>.

- Shum, M., Kelsh, M.A., Sheppard, A.R., Zhao, K., 2011. An evaluation of self-reported mobile phone use compared to billing records among a group of engineers and scientists. *Bioelectromagnetics* 32, 37–48. <https://doi.org/10.1002/bem.20613>.
- Smart, M., Pettis, J., Rice, N., Browning, Z., Spivak, M., 2016. Linking measures of colony and individual honey bee health to survival among apiaries exposed to varying agricultural land use. *PLoS One* 11, e0152685. <https://doi.org/10.1371/journal.pone.0152685>.
- Stange, E., Zulian, G., Rusch, G., Barton, D., Nowell, M., 2017. Ecosystem services mapping for municipal policy: ESTIMAP and zoning for urban beekeeping. *One Ecosyst.* 2, e14014. <https://doi.org/10.3897/oneeco.2.e14014>.
- Steinhauer, N., Kulhanek, K., Antúnez, K., Human, H., Chantawannakul, P., Chauzat, M.-P., VanEngelsdorp, D., 2018. Drivers of colony losses. *Curr. Opin. Insect Sci.* 26, 142–148. <https://doi.org/10.1016/j.cois.2018.02.004>.
- Straub, L., Williams, G.R., Pettis, J., Fries, I., Neumann, P., 2015. Superorganism resilience: eusociality and susceptibility of ecosystem service providing insects to stressors. *Curr. Opin. Insect Sci.* 12, 109–112. <https://doi.org/10.1016/j.cois.2015.10.010>.
- Urbiniello, D., Joseph, W., Huss, A., Verloock, L., Beekhuizen, J., Vermeulen, R., Martens, L., Röösl, M., 2014a. Radio-frequency electromagnetic field (RF-EMF) exposure levels in different European outdoor urban environments in comparison with regulatory limits. *Environ. Int.* 68, 49–54. <https://doi.org/10.1016/j.envint.2014.03.007>.
- Urbiniello, D., Joseph, W., Verloock, L., Martens, L., Röösl, M., 2014b. Temporal trends of radio-frequency electromagnetic field (RF-EMF) exposure in everyday environments across European cities. *Environ. Res.* 134, 134–142. <https://doi.org/10.1016/j.envres.2014.07.003>.
- Vacha, M., Puzova, T., Kvalcova, M., 2009. Radio frequency magnetic fields disrupt magnetoreception in American cockroach. *J. Exp. Biol.* 212, 3473–3477. <https://doi.org/10.1242/jeb.028670>.
- vanEngelsdorp, D., Meixner, M.D., 2009. A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *J. Invertebr. Pathol.* 103, S80–S95. <https://doi.org/10.1016/j.jip.2009.06.011>.
- vanEngelsdorp, D., Hayes, J., Underwood, R.M., Pettis, J., 2008. A survey of honey bee colony losses in the U.S., fall 2007 to spring 2008. *PLoS One* 3, 8–13. <https://doi.org/10.1371/journal.pone.0004071>.
- vanEngelsdorp, D., Evans, J.D., Saegerman, C., Mullin, C., Haubruge, E., Nguyen, B.K., Frazier, M., Frazier, J., Cox-Foster, D., Chen, Y., Underwood, R., Tarpy, D.R., Pettis, J.S., 2009. Colony collapse disorder: a descriptive study. *PLoS One* 4, e6481. <https://doi.org/10.1371/journal.pone.0006481>.
- Vermeeren, G., Markakis, I., Goeminne, F., Samaras, T., Martens, L., Joseph, W., 2013. Spatial and temporal RF electromagnetic field exposure of children and adults in indoor micro environments in Belgium and Greece. *Prog. Biophys. Mol. Biol.* 113, 254–263. <https://doi.org/10.1016/j.pbiomolbio.2013.07.002>.
- Verschaeve, L., 2014. Environmental impact of radiofrequency fields from mobile phone base stations. *Crit. Rev. Environ. Sci. Technol.* 44, 1313–1369. <https://doi.org/10.1080/10643389.2013.781935>.
- Vilić, M., Tlak Gajger, I., Tucak, P., Štambuk, A., Šrut, M., Klobučar, G., Malarić, K., Žura Žaja, I., Pavelić, A., Manger, M., Tkalec, M., 2017. Effects of short-term exposure to mobile phone radiofrequency (900 MHz) on the oxidative response and genotoxicity in honey bee larvae. *J. Apic. Res.* 56, 430–438. <https://doi.org/10.1080/00218839.2017.1329798>.
- Waldmann-Selsam, C., Balmori-de la Puente, A., Breunig, H., Balmori, A., 2016. Radiofrequency radiation injures trees around mobile phone base stations. *Sci. Total Environ.* 572, 554–569. <https://doi.org/10.1016/j.scitotenv.2016.08.045>.
- Wallaczek, J., 1992. Electromagnetic field effects on cells of the immune system: the role of calcium signalling. *Faseb J.* 6, 3177–3185. <https://doi.org/10.1080/14427591.2010.9686677>.
- Wang, Y., Ma, L.-T., Xu, B.-H., 2015. Diversity in life history of queen and worker honey bees, *Apis mellifera* L. *J. Asia Pac. Entomol.* 18, 145–149. <https://doi.org/10.1016/j.aspen.2014.11.005>.
- Weisbrot, D., Lin, H., Ye, L., Blank, M., Goodman, R., 2003. Effects of mobile phone radiation on reproduction and development in *Drosophila melanogaster*. *J. Cell. Biochem.* 89, 48–55.
- World Health Organization WHO, 2014. Electromagnetic fields and public health: mobile phones. <http://www.who.int>, Accessed date: 16 August 2018.
- Youngsteadt, E., Appler, R.H., López-Urbe, M.M., Tarpy, D.R., Frank, S.D., 2015. Urbanization increases pathogen pressure on feral and managed honey bees. *PLoS One* <https://doi.org/10.1371/journal.pone.0142031>.